



Published in final edited form as:

*Dis Colon Rectum*. 2010 August ; 53(8): 1148–1154. doi:10.1007/DCR.0b013e3181d5da6c.

## Factors Impacting quality of life in Women with fecal incontinence

Alayne D. Markland, DO, MSc<sup>1,2</sup>, W. Jerod Greer, MD<sup>3,4</sup>, Alicia Vogt, MD<sup>4</sup>, David T. Redden, PhD<sup>1,5</sup>, Patricia S. Goode, MSN, MD<sup>1,2</sup>, Kathryn L. Burgio, PhD<sup>1,2</sup>, and Holly E. Richter, PhD, MD<sup>3,4</sup>

<sup>1</sup> Birmingham/Atlanta Geriatric Research, Education, and Clinical Center (GRECC) Department of Veterans Affairs Medical Center, Birmingham, AL

<sup>2</sup>University of Alabama at Birmingham, Department of Medicine, Division of Gerontology, Geriatrics and Palliative Care

<sup>3</sup>University of Alabama at Birmingham, Department of Obstetrics and Gynecology

<sup>4</sup>Division of Women's Pelvic Medicine and Reconstructive Surgery

<sup>5</sup>University of Alabama at Birmingham School of Public Health

### Abstract

**Objective**—To characterize differences in health-related quality of life among women presenting for treatment of fecal incontinence.

**Methods**—Among 155 women presenting for treatment of fecal incontinence in a specialty clinic, validated questionnaires measured impact on quality of life (Modified Manchester Health Questionnaire) and severity (the Fecal Incontinence Severity Index). Bowel symptoms, including frequency, urgency, and stool consistency, were ascertained. Co-morbid diseases were self-reported. Linear regression models were constructed from significant univariate variables to examine differences seen in quality of life scores.

**Results**—Average age was  $58.7 \pm 11.5$  with no differences found in quality of life scores according to race, body mass index, or number of vaginal deliveries ( $p > 0.05$ ). Younger age, increased urinary incontinence symptoms, prior cholecystectomy, prior hysterectomy, and severity of bowel symptoms correlated with a negative impact on quality of life in univariate analysis ( $p$

---

Corresponding Author: Alayne D. Markland, DO, MSc, VA Medical Center, GRECC/11G, 700 19<sup>th</sup> St S., Birmingham, AL 35233, Phone: 205/934-3259, FAX: 205/558-7068, amarkland@aging.uab.edu.

Disclosures: None reported

#### Author Contributions:

- Alayne D. Markland, DO, MSc, Role: Planning and conducting study, interpreting data, drafting manuscript, Final draft approved by this author.
- W. Jerod Greer, MD, Role: Planning and interpreting data, drafting manuscript, Final draft approved by this author.
- Alicia Vogt, MD, Role: Data collection, interpreting data, and drafting manuscript, Final draft approved by this author.
- David T. Redden, PhD, Role: Interpretation of data, Final draft approved by this author.
- Patricia S. Goode, MSN, MD, Role: Planning and interpreting data, drafting manuscript., Final draft approved by this author.
- Kathryn L. Burgio, PhD, Role: Planning and interpreting data, drafting manuscript., Final draft approved by this author.
- Holly E. Richter, PhD, MD, Role: Planning and conducting study, data collection, interpreting data, drafting manuscript, Final draft approved by this author.

<0.05). Average severity scores were  $30.5 \pm 13.7$  with moderate correlation seen with increasing severity and quality of life scores ( $R_2 = 0.60$ ). After controlling for severity, women had increased quality of life scores with more bowel urgency (15 points, 95% CI 8.1, 21.2), harder stool consistency (10 points, 95% CI 3.8, 16.3), and prior hysterectomy (9 points, 95% CI 2.7, 15.4).

**Conclusion**—Bowel symptoms and having a prior hysterectomy had the greatest negative impact on quality of life in women seeking treatment for fecal incontinence. Targeting individualized treatments to improve bowel symptoms may improve quality of life for women with fecal incontinence.

### Keywords

fecal incontinence; women; quality of life; urinary incontinence; bowel symptoms

## Introduction

Fecal incontinence (FI) is defined as the involuntary loss of solid or liquid feces that is a social or hygienic problem.<sup>1</sup> When analyzing the prevalence of at least monthly FI among community-dwelling women in the US, rates vary from 3-22% with older women consistently experiencing more FI than younger age groups.<sup>2-8</sup> Given the projected 38% increase in women aged 65 and older among the US population in the next few decades, more information is needed regarding differences in women presenting for FI treatment and the impact on quality of life (QOL).<sup>9</sup>

FI has a significant negative impact on QOL among women,<sup>10-14</sup> with few studies examining factors which impact QOL in women presenting for treatment.<sup>13</sup> FI may be associated with prior trauma to the external sphincter (obstetrical or prior sphincter surgery), alterations in rectal capacity and compliance, bowel symptoms such as urgency and frequency, and age-related thinning in the external anal sphincter and other pelvic muscles. Medical conditions or co-morbidities, such as diabetes, vascular disease, obesity, and neurologic diseases, are also independently associated with FI.<sup>5-7, 10</sup> Existing studies have examined differences in women presenting for FI treatment, but few have controlled for medical conditions and for patient centered outcomes, such as severity, on QOL.<sup>13</sup>

Given the paucity of data on QOL differences seen in women with FI, this study provides additional data on factors that may impact QOL in women presenting for FI treatment in a specialty-based clinic. After controlling for patient age and the severity of FI, we hypothesized that women presenting for FI treatment would have a greater negative impact on condition-specific QOL if they had more medical co-morbidities, prior trauma to the external sphincter, and increased bowel symptoms.

## Methods

### Participants

Participants included women who presented with a primary complaint of fecal incontinence to the Genito-Rectal Disorders Clinic at the University of Alabama at Birmingham (UAB) since 2003. All women provided written consent and the UAB Institutional Review Board (IRB) approved all data collection.

### Questionnaires

Women were mailed questionnaires on fecal and urinary incontinence prior to presenting the Genito-Rectal Disorders Clinic. Demographic information, past medical history, and other potential risk factors for fecal incontinence were collected and entered into an IRB-approved

database at the time of initial presentation to the clinic. Questionnaires to assess FI included the Modified Manchester Health Questionnaire (MMHQ), which is a validated questionnaire that also includes the Fecal Incontinence Severity Index (FISI).<sup>15, 16</sup> The MMHQ measures health-related quality of life (HR-QOL) for FI and includes 8 subscales on the impact of FI on: overall impact, role limitations, physical/social limitations, personal relationships, emotions, sleep/energy, sexual activity, and lifestyle adaptation. The MMHQ is scaled from 0-100, for total and subscale scores, where higher scores represent a negative impact on HR-QOL. Women who reported not being currently sexual active did not respond to questions on the impact of FI on this domain, thus only 7 subscales were averaged for the calculation of the total MMHQ score in women who were not sexually active. The FISI measures the frequency of liquid, solid, mucus, or flatal incontinence that occurs from “two or more times per day,” “once per day,” “two or more times per week,” “once a week,” to “one to three times per month.” Patient-weighted scores are used to determine severity and scores range from 0-61, with higher scores indicating worse FI severity.<sup>16</sup> A FISI score of 0 indicates continence. Women who responded to any level of flatal incontinence without having incontinence to liquid, solid, or mucous stool were considered incontinent to flatus only and not considered to have fecal incontinence. Only women with incontinence to liquid, solid, or mucous stool were included in this study.

Additional information from the MMHQ that is not used for the total or subscale scores includes questions on bowel movement frequency, fecal urgency, and stool consistency (Table 1). Bowel movement frequency was ascertained by the question, “How often do you move your bowels each day?” Response levels were: “1-2 times a day,” “3-4 times a day,” “5-6 times a day,” “7 or more times a day,” “every other day,” and “less than every other day.” Frequency was then categorized as “≤ every other day,” “1-2 times a day,” and “≥3-4 times a day.” Fecal urgency was evaluated by the question, “How often do you have a strong desire to move your bowels which makes you rush to the toilet?” The response levels are: “never,” “occasionally,” “sometimes,” “most of the time,” and “all of the time.” We defined fecal urgency as occurring “most” or “all of the time” with a strong desire to have a bowel movement and non-urgency as “never,” “occasionally,” and “sometimes.” Stool consistency was characterized by the questions: “What percent of your bowel movements are hard or little balls?” and “What percent of your bowel movements are loose or watery?” Women responded to categories ranging from zero to one-hundred percent. Stool consistency was then categorized as occurring ≥ 50% or <50% for hard/little balls and for loose/watery stool.

A modified Sandvik Severity Index was used to describe urine leakage as present or absent.<sup>17</sup> The Medical, Epidemiologic, and Social Aspects of Aging Questionnaire (MESA) was used to evaluate symptoms of urinary incontinence (UI).<sup>18</sup> The MESA has two subscales; one for stress urinary incontinence symptoms and one for urge urinary incontinence symptoms. Questions referring to loss of urine at times of exertion such as laughing, sneezing, lifting, or bending define symptoms of stress urinary incontinence. Questions referring to urine loss preceded by an urge to void, or uncontrollable voiding with little or no warning define symptoms of urge incontinence. The questionnaire has a range of scores of 0-27 for stress UI symptoms and 0-18 for urge UI symptoms. Higher scores represent greater symptom severity.

Women also completed the SF-12, a generic health-related quality of life questionnaire.<sup>19</sup> For this analysis, the scales of the SF-12 included the mental and physical component summary scores. For the MMHQ, a higher score indicates worse symptom bother or a negative impact of symptoms on daily functioning.<sup>15</sup> For the scales of the SF-12, a higher score indicates a positive impact on health-related quality of life.

## Clinical Evaluation with Anorectal Manometry and Endoanal Ultrasound

Anorectal manometry was completed by physicians using a water-perfused disposable catheter system (Medtronic, Inc Minneapolis, MN). Pressures were recorded during resting, squeezing, and pushing at 1 cm intervals starting at 5-6 cm from the anal verge. Maximum tolerated rectal volume was measured in milliliters using an air-filled balloon. To evaluate for tears or disruption of the internal and external anal sphincters, endoanal ultrasound using a 10 MHz, 360° window endoanal probe at 5mm intervals (B & K Medical Systems, Inc. Willmington, MA) was completed. Women who had scarring of the external anal sphincter were also documented, but not included in the sphincter tear group.

## Data Analysis

Pearson's correlation coefficient was calculated between FISIQ scores and MMHQ scores. The relationship with all potential variables and the change in MMHQ scores was evaluated with one-way ANOVA, where a positive change represents an increase in MMHQ score (negative impact on quality of life) and a negative change representing a decline in MMHQ score (positive impact on quality of life). Data are presented as coefficients with 95% confidence intervals. In order to explore those factors that may impact differences in HR-QOL seen in women seeking treatment for FI, a step-wise linear regression model controlling for FI severity and co-morbid disorders with significant differences ( $P < 0.1$ ) on univariate analyses was constructed. Software used for analyses was STATA 8.2 (College Station, TX).

## Results

Our cohort included 164 women seeking evaluation for a primary complaint of FI. A subset of women (155/164, 95%) had complete data on the MMHQ. Age ranged from 21 to 85 years (mean age  $\pm$  standard deviation was  $58.7 \pm 11.5$  years). The sample was predominantly Non-Hispanic White participants (88%), with 9% African American, 2% Asian American, and 1% self-reported as "Other." Table 2 displays clinical characteristics of participants ( $n = 155$ ) and relationship with MMHQ scores. Younger women were more likely to have higher MMHQ scores (representing a negative impact on HR-QOL). When dichotomizing age, women  $< 65$  years of age had 15-point higher MMHQ scores than women  $\geq 65$  years of age. Other variables found to increase MMHQ scores ( $p \leq 0.05$ ) were: the presence of urinary incontinence (with increasing stress and urge symptoms), prior cholecystectomy, and prior hysterectomy.

In terms of QOL, mean SF-12 physical and mental scores were  $39.0 \pm 5.7$  and  $39.1 \pm 5.8$ , respectively, with a weak correlation seen with FISIQ scores ( $R_2 < -0.1$  for physical and mental subscale scores). Mean FISIQ scores were  $30.5 \pm 13.2$  and mean MMHQ scores were  $45.4 \pm 24.5$ . When examining MMHQ scores in comparison to FI severity, scores increased moderately ( $R_2 = 0.6$ ) with FISIQ scores. Only 51% of women aged  $\geq 65$  years of age with FI reported currently being sexual active at the time of the questionnaire compared to 80% of women  $< 65$  years of age (data not shown). Having more bowel symptoms including urgency and increased bowel movement frequency increased MMHQ scores by 25 points ( $p < 0.001$ ) and 7 points ( $p = 0.02$ ), respectively (Table 2). Stool consistency that was hard/little balls  $\geq 50\%$  of the time increased MMHQ scores by 14 points ( $p = 0.001$ ) and loose/watery stool consistency  $\geq 50\%$  of the time increased MMHQ scores 16 points ( $p < 0.001$ ).

Anorectal manometry and endoanal ultrasound were performed in a subset of the women presenting for FI treatment, 77% and 86%, respectively (Table 2). No significant changes in MMHQ scores were seen when evaluating objective measures such as anorectal manometry pressures and capacity. Although women who had a tear on endoanal ultrasound in either the

external or internal anal sphincters had increased MMHQ scores, this relationship was not significant.

After controlling for age, co-morbid disease, and FI severity in the linear regression model (Table 3), women reporting strong urgency symptoms had a 15-point higher MMHQ score (95% Confidence Interval 8 to 21). Women with hard stools or little balls  $\geq 50\%$  of the time had a 10-point increase in MMHQ scores. In multivariable analysis, loose or watery stool consistency was not a significant factor for increased MMHQ scores ( $p = 0.09$ ) and had no interaction with urgency symptoms. Having a prior hysterectomy was associated with a 9-point increase in MMHQ scores (95% Confidence Interval 3 to 15). The decrease in MMHQ scores seen in older women remained significant after controlling for other factors. If age was dichotomized as  $\geq 65$  years of age versus  $< 65$  years of age in the multivariable model (data not shown), the decrease in MMHQ scores in women  $\geq 65$  years of age remained significant and had a larger magnitude of change (-9.4 points decline, 95% Confidence Interval -16.6 to -2.2). With age dichotomized in the multivariable model, having a prior hysterectomy no longer remained a significant factor impacting MMHQ scores (6.2-point increase, 95% Confidence Interval -0.1 to 12.4).

## Discussion

Even after controlling for age and co-morbid diseases in this clinical cohort of women presenting for FI treatment, we found that bowel urgency symptoms, hard stool consistency, and having had a prior hysterectomy remained significant independent predictors of impaired HR-QOL. Although anal sphincter injury resulting from vaginal delivery is viewed as an important factor with regards to FI severity and QOL,<sup>20</sup> we found that bowel urgency and hard stool consistency had a greater negative impact on HR-QOL and deserve more attention from both a future research standpoint, as well as clinically, with assessment and treatment of FI.

Other studies have also found that bowel urgency and stool consistency are associated with FI in community-dwelling women,<sup>2,11,21</sup> but very few studies have examined the effect of bowel symptoms on HR-QOL in women with FI.<sup>13</sup> In one study that evaluated the impact of FI on HR-QOL in 157 patients attending urogynecology and colorectal clinics for various problems, increased bowel frequency and the amount of soiling had a negative impact on the Fecal Incontinence Quality of Life (FIQL) Scale.<sup>13</sup> Other studies have evaluated the impact of bowel symptoms on the presence of FI. In one such study by Bharucha et al, rectal urgency and symptoms of incomplete evacuation were independent risk factors for FI using a mailed bowel symptoms questionnaire among community-dwelling women.<sup>21</sup> In addition to fecal urgency, stool consistency plays an important role in FI and may impact fecal urgency. National data from the National Health and Nutritional Examination Survey (NHANES) from 2005-2006 found that advancing age, loose or watery stools, increased stool frequency, multiple chronic illnesses, and urinary incontinence were independent risk factors for FI in women, but this study did not assess symptoms of rectal urgency or the impact of bowel symptoms on QOL.<sup>8</sup> Although we did not use a validated measure to assess stool consistency, we did address the impact of stool consistency on QOL as measured by MMHQ questions. These findings may have important implications when doing FI intervention trials on modifiable risk factors and evaluating outcomes associated with HR-QOL.

Our study adds to the current literature on HR-QOL in women with FI by further evaluating and controlling for co-morbid conditions and FI severity, which are independent risk factors for FI in other studies and may also negatively impact QOL. In one other study that looked at the impact of increased FI severity on HR-QOL on the FIQL, the authors found a weak



linear correlation ( $R_2 \leq 0.4$ ) with FI severity (FISI scores) and general QOL as measured by the SF-36 ( $R_2 \leq 0.1$ ).<sup>22</sup> The lack of correlation of the FISI scores and the FIQL may be due to the scoring of the FIQL into 4 different domains, whereas the MMHQ is scored from zero to one hundred for a total score, as well as the 8 subscales. In this study, we found a moderate correlation ( $R_2 = 0.6$ ) for increasing FI severity and a negative impact on HR-QOL. We also found a weak correlation of the SF-12 subscales with the MMHQ scores ( $R_2 \leq 0.1$ ). Other studies utilizing different measures of FI severity and HR-QOL have found that increasing FI frequency is associated with a negative impact on QOL.<sup>5, 23</sup> More exploration is needed for existing measures of FI severity and HR-QOL, especially in terms of the minimal important difference needed to power future intervention studies for FI treatment.

Interestingly, we found that hard stool consistency had a greater negative impact on HR-QOL than liquid stool consistency. As a limitation of this study, the evaluation of stool consistency was ascertained from a question on the MMHQ (Table 1) and not from the validated Bristol Stool Form Scale.<sup>24</sup> Others have noted that FI to liquid stool had a greater negative impact on general QOL (SF-36 scores) than solid stool FI.<sup>11</sup> But, these studies did not address overall stool consistency in terms of diarrhea or constipation, but looked at only the type FI as liquid or solid stool loss.<sup>11</sup> Severe constipation and FI were found to have the greatest negative impact on QOL in a study that compared QOL among benign anorectal disorders.<sup>25</sup>

Women who had a prior hysterectomy had a significant negative impact on HR-QOL. Unfortunately, we did not collect information on menopausal status or the use of hormones to explore the role of hormonal status related to prior hysterectomy and the differential impact of each on HR-QOL. One small randomized controlled trial looked at changes in FI symptoms after administration of estrogen cream and did not find an improvement in FI.<sup>26</sup> Others have evaluated the presence of estrogen receptors in the anorectal canal with mixed findings.<sup>27</sup> More data are needed to evaluate the impact of menopause and hormonal changes on FI in women.

Little is known about the age-related differences on HR-QOL in women with FI. In a study by Wang et al, older women with FI were more likely to have a negative impact on HR-QOL than the younger women, but these findings were not adjusted for obstetrical factors.<sup>12</sup> It is possible that women 65 years of age and older were less likely to report an overall negative impact on HR-QOL than the younger women due to treatment-seeking differences for FI. Younger women may have been more likely to report and seek treatment for their FI symptoms in this cohort. Older women may have been more likely to adapt to their FI symptoms with protective undergarments, as well as limiting social interactions and travel outside the home. The MMHQ subscales do try to address these adaptive behaviors for women with FI and no significant differences were seen according to age in the lifestyle adaptation subscale of this instrument (data not shown).

Our study has a few other limitations that deserve further discussion. The main limitations of our study lie in its retrospective nature and the potential for missing data. Although data were collected prospectively, missing data were obtained through a retrospective chart review. We also had limited obstetrical data for instrumented deliveries. Another limitation of this single site study was the makeup of our patient population being nearly 90% Caucasian and the potential for referral bias, limiting the generalizability of our results. Strengths of our study include the characterization of the patient population, the use of validated instruments to assess FI severity and HR-QOL, and physiological testing with anorectal manometry and endoanal ultrasound. We plan to continue to characterize these

patients after various treatment modalities to further understand factors that may impact on improvement in symptoms.

## Conclusions

Our study confirms findings of previous research, that bowel symptoms, especially urgency and stool consistency, are important factors when addressing patient-centered outcomes such as QOL for FI. Assessment and treatment of fecal urgency and stool consistency may be important factors to consider when measuring patient-centered outcomes in future FI treatment trials.

## Acknowledgments

The authors would like to acknowledge Teresa McCool, RN and Cherry McNeely for their assistance with the database.

**Funding:** K24DK068389-03 to Dr. Richter Veterans Health Administration Career Development Award (CDA-2) to Dr. Markland Hartford/Southeast Center of Excellence in Geriatric Medicine Grant to Dr. Markland

## References

1. Norton, C.; Whitehead, WE.; Bliss, DZ., et al. Conservative and pharmacologic management of fecal incontinence in adults. In: Abrams, P.; Cardozo, L.; Khoury, S.; Wein, A., editors. Incontinence, 3rd International Consultation on Incontinence. Paris, France: Health Publications Ltd; 2005. p. 1521-64.
2. Bharucha AE, Zinsmeister AR, Locke GR, Seide BM, McKeon K, Schleck CD, Melton LJ. Prevalence and burden of fecal incontinence: A population-based study in women. *Gastroenterology* 2005;105(1):42. [PubMed: 16012933]
3. Macmillan AK, Merrie AEH, Marshall RJ, Parry BR. The Prevalence of Fecal Incontinence in Community-Dwelling Adults: A Systematic Review of the Literature. *Dis Colon Rectum* 2004;47:1341-9. [PubMed: 15484348]
4. Nelson R, Norton N, Cautley E, et al. Community-based prevalence of anal incontinence. *JAMA* 1995;274:559-561. [PubMed: 7629985]
5. Varma M, Brown J, Creasman J, Thom DH, Van Den Eeden SK, Beattie MS, Subak LL. Reproductive Risks for Incontinence Study at Kaiser (RRISK) Research Group. Fecal Incontinence in Females Older Than Aged 40 Years: Who is at Risk? *Dis Colon Rectum* 2006;49:841-51. [PubMed: 16741640]
6. Melville JL, Fan MY, Newton K, Fenner D. Fecal incontinence in US women: a population-based study. *Am J Obstet Gynecol* 2005;193:2071-76. [PubMed: 16325618]
7. Goode PS, Burgio KL, Halli AD, Jones RW, Richter HE, Redden DT, Baker PS, Allman RM. Prevalence and correlates of fecal incontinence in community-dwelling older adults. *J Am Geriatr Soc* 2005;53:629-35. [PubMed: 15817009]
8. Whitehead WE, Borrud L, Goode PS, Meikle S, Mueller ER, Tuteja A, Weidner A, Weinstein M, Ye W. Pelvic Floor Disorders Network. Fecal incontinence in US Adults: Epidemiology and risk factors. *Gastroenterology* 2009;137:512-7.
9. Administration on aging: Profile of older americans 2008. [25 March, 2009]. Available at: <http://www.aoa.gov/prof/Statistics/profile/profiles.aspx>;
10. Quander CR, Morris MC, Melson J, Bienias JL, Evans DA. Prevalence of and factors associated with fecal incontinence in a large community study of older individuals. *Am J Gastroenterol* 2005;100:905-09. [PubMed: 15784039]
11. Boreham MK, Richter HE, Kenton KS, Nager CW, Gregory WT, Aronson MP, Vogt VY, McIntire DD, Schaffer JI. Anal incontinence in women presenting for gynecologic care: prevalence, risk factors, and impact upon quality of life. *Am J Obstet Gynecol* 2005;192:1637-42. [PubMed: 15902170]

12. Wang JY, Patterson TR, Hart SL, Varma MG. Fecal incontinence: does age matter? Characteristics of older vs younger women presenting for treatment of fecal incontinence. *Dis Colon Rectum* 2008;51:426–31. [PubMed: 18213493]
13. Bartlett L, Nowak M, Ho Y. Impact of fecal incontinence on quality of life. *World J Gastroenterol* 2009;15:3276–82. [PubMed: 19598304]
14. Markland AD, Richter HE, Kenton KS, Wai C, Nager CW, Kraus SR, Xu Y, Tennstedt SL, for the Urinary Incontinence Treatment Network. Associated Factors and Impact of Fecal Incontinence in Women with Urge Urinary Incontinence. *Am J Obstet Gynecol* 2009;566:e1–6.
15. Kwon S, Visco AG, Fitzgerald MP, Ye W, Whitehead WE. Validity and Reliability of the Modified Manchester Health Questionnaire in Assessing Patients With Fecal Incontinence. *Dis Colon Rectum* 2005;48(2):323. [PubMed: 15616750]
16. Rockwood TH, Church JM, Fleshman JW, Kane RL, Mavrantonis C, Thorson AG, Wexner SD, Bliss D, Lowry AC. Patient and surgeon ranking of the severity of symptoms associated with fecal incontinence: the fecal incontinence severity index. *Dis Colon Rectum* 1999;42:1525–32. [PubMed: 10613469]
17. Sandvik H, Seim A, Vanvik A, Hunskaar S. A severity index for epidemiological surveys of female urinary incontinence: comparison with 48-hour pad-weighing tests. *Neurourol Urodyn* 2000;19(2):137–45. [PubMed: 10679830]
18. Diokno AC, Brock BM, Brown MB, Herzog AR. Prevalence of urinary incontinence and other urological symptoms in the noninstitutionalized elderly. *J Urol* 1986;136:1022–25. [PubMed: 3490584]
19. Ware JE, Kosinski M, Keller SD. A 12-Item Short-Form Health Survey: Construction of scales and preliminary tests of reliability and validity. *Medical Care* 1996;34:220–33. [PubMed: 8628042]
20. Madoff RD, Parker SC, Varma MG, Lowry AC. Faecal incontinence in adults. *Lancet* 2004;364:621–32. [PubMed: 15313364]
21. Bharucha AE, Zinsmeister AR, Locke GR, Schlek C, McKeon K, Melton LJ. Relationship of bowel habits to fecal incontinence in women. *Clin Gastroenterol Hepatol* 2006;44:1004–9. [PubMed: 16630766]
22. Bordeianou L, Rockwood T, Baxter N, Lowry AC, Mellgren A, Parker S. Does incontinence severity correlate with quality of life? *Colorectal Dis* 2008;10(3):273–79.
23. Baxter NN, Rothenberger DA, Lowry AC. Measuring fecal incontinence. *Dis Colon Rectum* 2003;46:1591–605. [PubMed: 14668583]
24. Heaton KW, Radvan J, Cripps H, Mountford RA, Braddon FE, Hughes AO. Defecation frequency and timing, and stool form in the general population: a prospective study. *Gut* 1992;33:818–24. [PubMed: 1624166]
25. Sailer M, Bussen D, Debus ES, Fukes KH, Thiede A. Quality of life of patients with benign anorectal disorders. *B J Surg* 1998;85:1716–9.
26. Pinedo G, García E, Zárate AJ, León F, Bellolio F, Molina ME, Viviani P, Á Z. Are topical oestrogens useful in faecal incontinence? Double-blind randomized trial. *Colorectal Dis* 2009;11:390–3. [PubMed: 18637100]
27. Rizk DE, Helal TE, Mason N, Berg B. Non-evidence of estrogen receptors in the rectal mucosa. *Int Urogynecol Pelvic Floor Dysfunct J* 1998;9:136–9.



**Table 1**

## Bowel Symptom Questions and Responses from the Modified Manchester Health Questionnaire

Question	Bowel symptoms	Response levels	N (%) of women responding
“How often do you move your bowels each day?”	Frequency	• 1-2 times/day	65 (46%)
		• 3-4 times/day	45 (31%)
		• 5-6 times/day	6 (4%)
		• 7 or more times/day	3 (2%)
		• every other day	10 (7%)
		• < every other day	14 (10%)
“How often do you have a strong desire to move your bowels which makes you rush to the toilet?”	Urgency	• never	9 (6%)
		• occasionally	23 (16%)
		• sometimes	59 (40%)
		• most of the time	44 (30%)
		• all of the time	12 (8%)
“What percent of your bowel movements are hard or little balls?”	Hard stool consistency	• 0%	36 (25%)
		• 25%	64 (43%)
		• 50%	22 (15%)
		• 75%	19 (13%)
		• 100%	6 (4%)
“What percent of your bowel movements are loose or watery?”	Loose stool consistency	• 0%	26 (18%)
		• 25%	48 (33%)
		• 50%	43 (29%)
		• 75%	25 (17%)
		• 100%	5 (3%)

**Table 2**

Univariate Analysis of Demographic Variables and Co-morbid Diseases Associated with Increased HR-QOL in Women Seeking Treatment for FI, n = 155

<i>Variables</i>	<i>N (%) or Mean ± standard deviation</i>	<i>MMHQ Score<sup>a</sup> Coefficient, (95% confidence interval)</i>
Age (years)	54 ± 14	-0.3 (-0.6, 0.0)
Age ≥ 65 years	43 (28%)	-15.0 (-23.26, -6.7)
Non-White race/ethnicity	16 (10%)	1.4 (-11.5, 14.3)
Body mass index (kg/m <sup>2</sup> )	27 ± 8	0.1 (-0.4, 0.5)
Diabetes	16 (10%)	3.3 (-9.6, 16.2)
Coronary artery disease	19 (12%)	-3.8 (-16.0, 8.5)
Hypertension	55 (35%)	3.9 (-4.4, 12.3)
Inflammatory bowel disease	16 (10%)	10.6 (-2.2, 23.3)
Urinary incontinence	69 (45%)	7.7 (-0.2, 15.6)
MESA stress scores	9 ± 6	0.6 (0.1, 1.1)
MESA urge scores	10 ± 8	0.9 (0.3, 1.6)
SF-12 subscales		
Physical	39 ± 6	-0.1 (-0.8, 0.6)
Mental	39 ± 6	-0.1 (-0.8, 0.6)
Cholecystectomy	35 (23%)	10.2 (0.8, 19.6)
Anorectal surgery (hemorrhoidectomy)	32 (21%)	-0.2 (-10.1, 9.8)
Cancer (other than skin)	14 (9%)	4.2 (-10.4, 18.9)
Hysterectomy	99 (64%)	8.7 (0.6, 16.8)
Prior c-section	14 (9%)	6.6 (-7.0, 20.3)
Number of vaginal deliveries	2 ± 1	0.7 (-2.3, 3.6)
<i>Bowel Symptoms Variables (n = 147, except FISI scores)</i>		
FISI scores (FI severity), n = 155	31 ± 13	1.1 (0.9, 1.3)
Fecal urgency symptoms, occurring "most" or "all of the time"	56 (38%)	24.5 (17.3, 31.8)
≥ 3-4 bowel movements/day	57 (39%)	6.8 (1.2, 12.3)
Bowel consistency hard or little balls, ≥ 50% of the time	47 (32%)	13.7 (5.4, 22.0)
Bowel consistency loose or watery, ≥ 50% of the time	73 (50%)	16.0 (8.4, 23.6)
<i>Anorectal Manometry (n = 127) and Endoanal Ultrasound Variables (n = 141)</i>		
Resting pressure (mmH <sub>2</sub> O)	34 ± 22	-0.1 (-0.3, 0.1)
Squeeze pressure (mmH <sub>2</sub> O)	64 ± 25	-0.1 (-0.2, 0.0)
Rectal capacity (ml)	138 ± 68	-0.0 (-0.1, 0.0)
External anal sphincter tear	60 (42%)	3.9 (-4.2, 12.0)
Internal anal sphincter tear	21 (15%)	8.4 (-3.0, 19.8)
Tear in both sphincters	16 (11%)	10.2 (-4.3, 24.6)

<sup>a</sup>MMHQ Scores vary from 0-100 with higher scores indicating a greater impact on HR-QOL. Positive coefficients represent increased MMHQ scores (negative impact on health-related quality of life); negative coefficients represent a decline in MMHQ scores (positive impact on health-related quality of life).

**Table 3**

Linear Regression Model of Variables Associated with Increased Impact on HR-QOL in Women Seeking Treatment for FI, n = 142

<i>Variables</i>	<b>MMHQ Score, Coefficients with 95% CI<sup>a</sup></b>		
	<b>CoE</b>	<b>95% CI</b>	<b>p value</b>
Urgency sensation	14.7	8.1, 21.2	<0.001
Stool consistency (hard or little balls) $\geq$ 50% of the time	10.0	3.8, 16.3	0.002
Age (years)	-0.3	-0.5, -0.1	0.01
Fecal incontinence severity	0.8	0.6, 1.1	<0.001
Hysterectomy	9.1	2.7, 15.4	0.005

<sup>a</sup>Model also controlled for the following variables: age, presence of urinary incontinence, prior cholecystectomy, bowel movement frequency, and loose stool consistency.